

Non-technical soil descriptions for Summit Soil Survey

101-Agassiz-rock outcrop complex, 30 to 70 percent slopes.

Agassiz, about 60 percent, is a shallow, somewhat excessively drained, moderately permeable soil on mountain slopes under sagebrush and grasses. Typically it is neutral very cobbly loam over limestone bedrock at 10 to 20 inches. Precipitation is 16 to 22 inches, frost-free period is 60 to 90 days, and capability rating is 7e nonirrigated.

Rock outcrop, about 20 percent, is on ridges and escarpments.

102—Ant Flat loam, 2 to 8 percent slopes is a very deep, well drained, slowly permeable soil on fan remnants under sagebrush and grasses or cropland. Typically it is neutral loam about 13 inches thick over clay loam or clay about 17 inches thick over moderately alkaline clay loam to 60 inches or more. Precipitation is 16 to 22 inches, frost-free period is 60 to 90 days, and capability rating is 3e irrigated and nonirrigated.

103—Ant Flat loam, 8 to 15 percent slopes is a very deep, well drained, slowly permeable soil on fan remnants under sagebrush and grasses or cropland. Typically it is neutral loam about 13 inches thick over clay loam or clay about 17 inches thick over moderately alkaline clay loam to 60 inches or more. Precipitation is 16 to 22 inches, frost-free period is 60 to 90 days, and capability rating is 4e irrigated and nonirrigated.

104—Ant Flat loam, 15 to 30 percent slopes is a very deep, well drained, slowly permeable soil on fan remnants under sagebrush and grasses. Typically it is neutral loam about 13 inches thick over clay loam or clay about 17 inches thick over moderately alkaline clay loam to 60 inches or more. Precipitation is 16 to 22 inches, frost-free period is 60 to 90 days, and capability rating is 6e nonirrigated.

105—Ant Flat-Henefer-Skutum complex, 8 to 30 percent slopes.

Ant Flat, about 40 percent, is a very deep, well drained, slowly permeable soil on mountain slopes under sagebrush and grasses. Typically it is neutral loam about 13 inches thick over clay loam or clay about 17 inches thick over moderately alkaline clay loam to 60 inches or more. Precipitation is 16 to 22 inches, frost-free period is 60 to 90 days, and capability rating is 6e nonirrigated.

Henefer, about 30 percent, is a very deep, well drained, slowly permeable soil on mountain slopes under oak brush. Typically it is neutral gravelly loam about 12 inches thick over cobbly clay or clay loam about 18 inches thick over neutral or slightly alkaline very cobbly or gravelly sandy clay loam to 60 inches or more. Precipitation is 16 to 22 inches, frost-free period is 60 to 90 days, and capability rating is 6e nonirrigated.

Skutum, about 20 percent, is a deep, well drained, slowly permeable soil on mountain slopes under aspen. Typically it is neutral loam about 17 inches thick over gravelly clay or clay loam about 31 inches thick over gravelly sandy loam over sandstone bedrock at 40 to 60 inches. Precipitation is 22 to 35 inches, frost-free period is 20 to 60 days, and capability rating is 6e nonirrigated.

permeable soil on mountain slopes under sagebrush and grasses. Typically it is slightly acid cobbly loam about 6 inches thick over slightly acid or neutral gravelly clay loam about 17 inches thick over neutral very cobbly loam over andesite bedrock at 20 to 40 inches. Precipitation is 16 to 22 inches, frost-free period is 60 to 90 days, and capability rating is 6s nonirrigated.

107-Ayoub-Dunferd-Melling complex, 15 to 30 percent slopes.

Ayoub, about 40 percent, is a moderately deep, well drained, moderately slowly permeable soil on mountain slopes under sagebrush and grasses. Typically it is slightly acid cobbly loam about 6 inches thick over slightly acid to neutral gravelly clay loam about 17 inches thick over neutral very cobbly loam over andesite bedrock at 20 to 40 inches. Precipitation is 16 to 22 inches, frost-free period is 60 to 90 days, and capability rating is 6e nonirrigated.

Dunford, about 20 percent, is a moderately deep, well drained, moderately slowly permeable soil on mountain slopes under oak brush. Typically it is neutral cobbly loam about 10 inches thick over gravelly clay loam over andesite bedrock at 20 to 40 inches. Precipitation is 16 to 22 inches, frost-free period is 60

to 90 days, and capability rating is 6e nonirrigated.

Melling, about 20 percent, is a shallow, well drained, moderately slowly permeable soil on mountain slopes under sagebrush and grasses. Typically it is neutral extremely stony loam over very cobbly clay loam over andesite bedrock at 10 to 20 inches. Precipitation is 16 to 22 inches, frost-free period is 60 to 90 days, and capability rating is 7s nonirrigated.

-Ayoub-Dunford-Melling complex, 30 to 60 percent slopes.


Ayoub, about 40 percent, is a moderately deep, well drained, moderately slowly permeable soil on mountain slopes under sagebrush and grasses. Typically it is slightly acid cobbly loam about 6 inches thick over slightly acid to neutral gravelly clay loam about 17 inches thick over neutral very cobbly loam over andesite bedrock at 20 to 40 inches. Precipitation is 16 to 22 inches, frost-free period is 60 to 90 days, and capability rating is 7e nonirrigated.

Dunford, about 20 percent, is a moderately deep, well drained, moderately slowly permeable soil on mountain slopes under oak brush. Typically it is neutral cobbly loam about 10 inches thick over gravelly clay loam over andesite bedrock at 20 to 40 inches. Precipitation is 16 to 22 inches, frost-free period is 60

to 90 days, and capability rating is 7e nonirrigated.

Melling, about 20 percent, is a shallow, well drained, moderately slowly permeable soil on mountain slopes under sagebrush and grasses. Typically it is neutral extremely stony loam over very cobbly clay loam over andesite bedrock at 10 to 20 inches. Precipitation is 16 to 22 inches, frost-free period is 60 to 90 days, and capability rating is 7e nonirrigated.

- 109—Cluff loam, 8 to 30 percent slopes, is a deep, well drained, slowly permeable soil on mountain slopes under mixed conifer. Typically it is slightly acid loam over gravelly loam about 9 inches thick over very gravelly loam about 7 inches thick over very cobbly clay loam about 4 inches thick over very cobbly clay over conglomerate bedrock at 40 to 60 inches. Precipitation is 22 to 35 inches, frost-free period is 20 to 60 days, and capability rating is 6e nonirrigated.
- 110—Cluff loam, 30 to 60 percent slopes, is a deep, well drained, slowly permeable soil on mountain slopes under mixed conifer. Typically it is slightly acid loam over gravelly loam about 9 inches thick over very gravelly loam about 7 inches thick over very cobbly clay loam about 4 inches thick over very cobbly clay over conglomerate bedrock at 40 to 60 inches. Precipitation is 22 to 35 inches, frost-free period is 20 to 60 days, and capability rating is 7e nonirrigated.
- 111—Crandall gravelly loam, 2 to 8 percent slopes, is a deep, well drained, moderately slowly permeable soil on mountain slopes under grasses and forbs. Typically it is neutral gravelly loam about 14 inches thick over very cobbly clay loam over slightly alkaline very cobbly loam over conglomerate bedrock at 40 to 60 inches. Precipitation is 22 to 35 inches, frost-free period is 20 to 60 days, and capability rating is 6s nonirrigated.





## Soil Test Report and

## Fertilizer Recommendations

**USU Analytical Labs** 

Appendix 1

Utah State University Logan, Utah 84322-4830 (435) 797-2217 (435) 797-2117 (FAX)

Date Received:
Date Completed:

10/13/98 10/16/98

Name: Address: Lon Thomas American Stone 4040 S 300 W

Salt Lake City UT 84107

County: Salt Lake

Lab Number:

98011512

Grower's Comments:

Acres in Field:

Identification:

Peoa

Crop to be Grown: Dryland Pasture

Soil	Test R	esults	Interpretations	Recommendations
Texture		Loam		
Lime		+	Normal	
pН		7.2	Normal	
Salinity - ECe mml	nos/cm	1.3		
Phosphorus - P	ppm	23		0 lbs P2O5/A
Potassium - K	ppm	269		0 lbs K2O/A
Nitrate-Nitrogen - N	ppm	•		40-60 lbs N/A
Zinc - Zn	ppm			
Iron - Fe	ppm			
Copper - Cu	ppm			
Manganese - Mn	ppm			
Sulfate-Sulfur - S	ppm	•		
SAR		1.22	Soil Not Sodic	
Organic Matter	%	3.57		

Notes

> 2mm = 14.0%

Total Nitrogen = 0.155 %

Nested Frequency of the Peoa Quarry (above)

Transact Location: Peca quarry cabove		í		į		•	Š	į	•	ğ	ខ្លួ	Ž	į	Š		,	É	į	CHANNAL SIZE: 1- SQUASQUI, Z- ZSQUAZSQUI, S- ZSQUAZSUSII, 4-SUQUIASSQUI					
	Number of Qu	Š	of Out	adrats	8																			
Oftent Original							σ	Quadrate	#									Frage	Frequency Summary by Plot Size	mmary	by Plot	.5.		
THE SPOKES								ļ	i										2		3		4	
	2	3	7	5 6	2	8	9 1	10 11	12	13	7	1.5	16 1	17 1	18 19	9 20	至	%Freq	Z.	%Freq	壬	%Fred	SE SE	%Freq
Epilobium brachycarpum 2	2		-			┢	-	<u> </u>	L		T	$\vdash$	$\vdash$	$\vdash$	$\vdash$	L		0	2	2	7	2	2	5
Polygonum douglasii 2	2	2	8	7	-	77	2	2 2	2	-	7	7	-	2	2	2	2	=	15 18	8	19	88	9	8
Bromus tectorum 2	2	2	2	2 3	7	ম	2	2	7	7	77	N	2	+	F	2	7	Ľ	5	88	188	8	82	8
Poa bulbosa	F	2	4	2	7	7	2	1 2		~	2	7	2	77	-	2	3	3 15	9	8	17	85	180	8
Rosa woodsii	2	2	2 4	1		-	F	2 4	N	₹	<b>†</b>	₹	-	4	Ë	2		0	0	g		88	22	8
Bromus japonicus	7	63	2	2		$\vdash$	$\vdash$	2	ल		$\vdash$	-	-	┝	-	_	Ľ	0		20		8	~	8
Outierrezia sarothrae	2	7	-	3 2	ন	4	6	L	ಣ		1	H	~	┝	۲	7	2	3	-	35	2	જ	12	8
Verbena bracteata		4	-	L		H	-	L	L		1	$\vdash$	┞	┞	1	L		8	0		L		-	3
Arabis selbyi		T	87		4	H	-	L			H	-	ļ.	H	$\vdash$	L	L	8	0	ľ	-	2	m	15
Artemisia spicata		H	F	7	7	ಣ	+	2			4	~	H	-	60	-		10	9	8	8	3	F	188
Artemisia triclentata		1	$\vdash$	L	T	十	ŀ	4			T	H	╀	╀	+	L			0	L			~	12
Orobiranche conymbosa		T	-	L	T	62	-	L			t	H	╀	┝	-	L		0	0				-	15
Alyssum alyssloides		T	-		T	$\vdash$	+	-	L	N	t	t	╀	╀	$\vdash$	L		0	10		-	*	F	l
Elymus spicatus		H	$\vdash$		T	$\vdash$	┞	-		4	=	2	┞	╀	+	Ļ			5	٦	2	2	6	15
Mahonia repens			-		T		-	-			T	~	6	2	-	L		0	2				F	15
Viguiera multiflora	Ħ	H	H	П	H	Н	Н	Н			H	H	H	$\vdash$	H	4		0	0				-	5
Covor Summary	Vegetation: 18%	¥	<b>%</b>		Litter	Litter: 40%	yg.		89	5	Bare Ground: 31%	31			<u> 8</u>	i de	Grawel/Strane: 13%	13%						

## Soil Test Report and

## Fertilizer Recommendations

Date Received:

10/13/98

Date Completed:

10/16/98

Name:

Lon Thomas

Address:

American Stone 4040 S 300 W

Salt Lake City UT 84107

County: Salt Lake

**USU Analytical Labs** 

**Utah State University** 

(435) 797-2117 (FAX)

(435) 797-2217

Logan, Utah 84322-4830

Lab Number:

98011512

Grower's Comments:

Acres in Field:

DIV. OF OIL, GAS & MINING

Identification:

Peoa

Crop to be Grown: Dryland Pasture

Soil Test Res	ults	Interpretations	Recommendations
Texture	Loam		
Lime	+	Normal	
рН	7.2	Normal	
Salinity - ECe mmhos/cm	1.3		
Phosphorus - P ppm	23		0 lbs P2O5/A
Potassium - K ppm	269		0 lbs K2O/A
Nitrate-Nitrogen - N ppm			40-60 lbs N/A
Zinc - Zn ppm			
Iron - Fe ppm			
Copper - Cu ppm			
Manganese - Mn ppm			
Sulfate-Sulfur - S ppm			
SAR	1.22	Soil Not Sodic	
Organic Matter %	3.57		

Notes

> 2mm = 14.0%

Total Nitrogen = 0.155 %

,				Allota		Number			
- DE	ile	Examine		7000					}
(case)		z	umber of Qua	drats				Qu	Quadrat Size
	Quadrat	S		ĺ	Fre	quency Sur	nmary by	Plot Size	
4 5 6	8 9 10	12 13	15 16 17	19	Hits %	His	Tiss	T	% Freq
								_	
3 2	1 2 2 2	2212	2 1 2	2 2					
2 3	222	/ अ	2 2	2 2				-	
N	N	2 2 2	2 2 2	123					
4	83	424	h h	2					
<b>9</b> 2 -		23						-	
ري د		2-7	2	2 2					
	4		4						
2	4		12 1						
	4	h .						_	
	L.							_	
		2						_	-
		4	2					-	
			233						
				4					-
							_		
	00000000								-
	N N N L N N 5	Date  Ouadrat  Quadrat  Quadrat	Date Examiner  Quadrats  Quadrats  2 2 1 2 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 2 1 2 2 2 1 2 2 2 1 2 2 2 1 2 2 2 1 2 2 2 1 2 2 2 1 2 2 2 1 2 2 2 2 1 2 2 2 2 1 2 2 2 2 1 2 2 2 2 1 2 2 2 2 1 2 2 2 2 1 2	Date Examiner  Quadrats  Quadrats	Date   Examiner   Number of Quadrats   Qua	Date   Examiner   Allotment Name   Country   Country	Date   Examiner   Allotment Name   Country   Country	Date   Examiner   Allotment Name   Country   Number of Quadrats   Family   Number of Quadrats	Date   Examiner   Allotment Name & Number   Vumber of Quadrats   Frequency Summary by Plot 8   9   10   11   12   13   14   15   16   17   18   19   20   Hs   % Freq   Hs   % Freq

Appendix 2.

				<b>Nested Frequency</b>	quency				Page of
Study Number		Date	Examiner		Allotment Name & Number	Number			Pasture
Transact I ocation			Numb	Number of Quadrats	20			Quadrat Size	
	1 mg / man	Quadrats				Frequency Summary by Plot Size	ary by Plot S	ize	Cover
Plant Species -	4 Э Э	A 7 8 9 10	11 12 13 14 15	5 16 17 18 19	20	- 1	Hits % Freq	Hits % Freq	Summary
_ _	4		2.	<b>1</b>		$\sim$			Vegetation (Basal)
there	-	300 XX			0	1 5	/   5	ر ا	<b>XX</b> 72%
in antien	)				$\dashv$	2 /0	2 /5	2 /<	ą
Ander rething			ST 5003		, 0	+	+		Hits %Cover
a lille mily ti	2	4 4	24	9 3	000	1/0	\\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	┵	Vegetati
	2 4	ca.	2	1 4	00	2/0	5 25	1	•
	2 4	2 3 2	34		00	3 /5	4 20	5 25	
and the state of t	7 2 1 2 2		/ 18 / 2	)	2 6	/3	14	15	Hite %Cover
Ostocarpin	9 4 2		9 3	(61	3 4 6		2	6	X X
instally.	(7	2			0			2	X M
	2324	4			<b>)</b>	5	3		484
licia					0	0	_		Hits %Cover
on	224	શ	H		0	2	ام ا	4	
idami	3 2	3 2	3 3 3		0	2	6	7	XX 799
25 7 07	1 1 1							) L.	
Sympacorpies					\ \	/	_	كر	Hits %Cover
repolition					0	0	/	R	Gravel/Stone
sto.		4 7 2	2 4			ر کا	3	6	
hysotlem,	446	2	4 3 3	ω 2	0	2	5	8	Hire %Cover
risoposo defe	3	2			0	~	<i>\</i>	)	Crytog
=	Offillering	2			0	_	_	_	
Oranium oran	tt		2		0	0	_	_	
where theres			- 1	record around	Dont record ground cover under car	nopy hits of shrubs under 10ft in height	ubs under 1	oft in height	Hits %Cover
res (Use off	(Use other side of allother payer)	l page/	(	W	0	0	~	_	

Nested Frequency of the Peoa Quarry (below)

Titherand   5 Part Ap. Done assure the bear		CAMPENDIA JAN	••	Spencer	₹		3	ğ	Quadrat Size	10	ž	5cm	7=2	Sea	S	3	: 25cm	.50cm:	ScnX5cm; 2= 25cmX25cm; 3= 25cmX50cm; 4=50cmX30cm	(30cm				
Tarbect Location: Pega ditarry (Delown		Number of		Quadrats:	4 1	20																		
								O	Quadrats								L	F	Frequency S	Summany	2	Plot Size		
Plant Species							į		į	i								-		7		3	_	
	-	2 3	4	5	9	7 8	6	<b>\$</b>	111	12	13 14	15	16	12	12	6	20 Hits	%Fired	ST K	%Frea	E E	%Fred	至	%Fred
Poa pretense	F	7		3	7	_		F	4	4	-	L		F	77	77	2	4	202	8	8	9	12	
Juncus articus	2			$\vdash$	_	L			H	$\vdash$	L	L		┪	H	┝	L			-	100	L	4	
Arabis selbyi	2	L		1	7	33			T	+	$\vdash$	L		T	t	╁	Ļ	0	-	N	2	3 15	3	5
Achillia millfolium	~			$\vdash$	-	4	4		T	+	2	-		4	H	╁	3	5	0	L	2	L		
Połygonum douglasii	2	1.4		H	೯	Ц			2	80	H	L			65	*	L	8	0	L	5	5 25	-	L
Epilobium brachycarpus		4		Н	2	3	2		H	2	Н	Ц				H		0	0	3	15	8	5	
Elymus spicata	60	2 1	2	2	_	4		2	+	-	7	2 1			F	2	2	9	30	13	88	14 70	15	25
Orthocarpus tolmei	+		4	2	Н				+	-	L	L		T	H	6	4	0	0	-	5	2 10	9	8
Dipsacus sylvestris	3			H	-	_	7		$\vdash$	-	L	L		T	H	H		6	0	-	9			
Artemisia tridentata	Ĥ	2 3	7	4	2	3	4	3	H	Н	7	21			က	┝	2	0	0	5	श्र	9 45		
Vicia americana	$\dashv$	+		$\vdash$	Н	3		П		Н	Н	L			H	-		0	0	0	0	1 5	2	
Rosa woodsii	$\dashv$	2	7	4		2		П	4	Н	Щ			一	-	H	L	0	8	6	55	3 15		
Mahonia repens	$\dashv$	ိ		7	$\dashv$	Ц	6	2	3	H	3 4	1		Н	Н	H		o	ō	7	2	9		L
Poa Builbosa	-	7		ᅱ	-	_				$\dashv$	Ц							1	S.	-	5	1 5	<u></u>	
synphorocarpus oreophilus	-		-	-	-	4				H	Ц			H	+	H	L	1	5	-	ψ.	1 5		15
Lipinus sparsiflorus	_			4	9			П			Н			Г	$\vdash$	$\vdash$	L	0	0	0	0	1 5	~	12
Aster chimensis	$\dashv$			Н	•	4 1	2		H	-	2 4	L		-	-	H	L	<u>-</u>	2	9	15	3 15	٥	
Chrysothamnus viscidifiorus	$\dashv$			Н	H	2		П	4	<del> </del>	3 3	3	4	-		┝	L	8	ō	2	9		8	
ragopogon dubius	4					2		-	┝	-	L	L		<del> </del>	-	┡	L	0	ø	-	5	1	_	3
Geranaum Viscocisimum	-			H	Н	Ц		2	H	$\vdash$	L					┞		0	0	-	5	1 5		5
Pursnia tridentata	4				_			П	H	Н	3	1		$\vdash$	$\vdash$	-		0	o	0	0	1 5		5
verbascum thaspus	4			$\dashv$	$\dashv$					$\vdash$		7			-	-	_	0	0	0	0	0	_	5
Amaranthus abus	4			$\dashv$	4			$\exists$	H	Н	Ц	3		П	Н	Н		0	0	0	8	1 5	-	3
Cover Summary	Vegetation: 23%	E E E	23%		_5	Litter: 48%	*		00	Bare Ground: 29%	, de	7 V	ž		Ľ	CO CO	Smartishne. (192	767						